V0401 ALCOHOL DETOX WITH DRUG DETOX PROCEDURES = COMBINATION

CODE NEEDED

Guideline: Detoxification is the management of withdrawal symptoms in a patient who is physically

dependent on alcohol or drugs. It is illogical to report drug detox and alcohol detox

separately when there is a combination code alcohol and drug detox.

V0401 Exclusive check (if match, error) - P017

Procedure Table 94.62 Alcohol detoxification

Relational Table 94.65 Drug detoxification

V0401 HINT: Instead of using two separate codes, report the combination code 94.68 Combined

alcohol and drug detoxification.

References: Coding Clinic for ICD-9-CM, AHA, 2nd Quarter 1991, pages 12-13.

Coding Clinic for ICD-9-CM, AHA, 3rd Quarter 1994, pages 13-14.

ICD-9-CM Coding Handbook With Answers, AHA, Faye Brown, RRA, 1991, pages 126-

128.

V0402 ALCOHOL REHAB WITH DRUG REHAB PROCEDURES = COMBINATION

CODE NEEDED

Guideline: Rehabilitation is a structured program carried out with the goal of establishing strict control

of the alcohol or drug. It is illogical to report drug rehab and alcohol rehab separately when

there is a combination code for alcohol and drug rehab.

V0402 Exclusive check (if match, error) - P018

Procedure Table 94.61 Alcohol rehabilitation

Relational Table 94.64 Drug rehabilitation

V0402 HINT: Instead of using two separate codes, report the combination code 94.67 Combined

alcohol and drug rehabilitation.

References: Coding Clinic for ICD-9-CM, AHA, 2nd Quarter 1991, pages 12-13.

Coding Clinic for ICD-9-CM, AHA, 3rd Quarter 1994, pages 13-14.

ICD-9-CM Coding Handbook With Answers, AHA, Faye Brown, RRA, 1991, pages 126-

128.

V0403 ALCOHOL REHAB and ALCOHOL DETOX PROCEDURES =

COMBINATION CODE NEEDED

V0404 DRUG REHAB and DRUG DETOX PROCEDURES =

COMBINATION CODE NEEDED

V0405 ALCOHOL/DRUG REHAB and DETOX PROCEDURES =

COMBINATION CODE NEEDED

Guideline:

When both detox and rehab are performed for both drug and alcohol, there is a combination code for combined alcohol and drug rehabilitation and detoxification. It is illogical to report combined alcohol and drug rehab separately from combined alcohol and drug detox when there is a combination code for combined alcohol and drug rehab and detox.

Multiple codes should not be assigned when the classification provides a combination code that clearly identifies all the elements documented in the procedure statement. Only the combination code is assigned when the code fully identifies the procedures involved.

V0403 Exclusive check (if match, error) - P019

Procedure Table 94.61 Alcohol rehabilitation

Relational Table 94.62 Alcohol detoxification

V0403 HINT: Instead of using two separate codes, report the combination code 94.63 Combined

alcohol rehabilitation and detoxification.

.....

V0404 Exclusive check (if match, error) - P020

Procedure Table 94.64 Drug rehabilitation

Relational Table 94.65 Drug detoxification

V0404 HINT: Instead of using two separate codes, report the combination code 94.66 Combined

drug rehabilitation and detoxification.

V0405 Exclusive check (if match, error) - P021

Procedure Table 94.67 Combined alcohol and drug rehabilitation

Relational Table 94.68 Combined alcohol and drug detoxification

V0405 HINT: Instead of using two separate codes, report the combination code 94.69 Combined

alcohol and drug rehabilitation and detoxification.

.....

References: Coding Clinic for ICD-9-CM, AHA, 2nd Quarter 1991, pages 12-13; 3rd Quarter 1994, pages

13-14.

ICD-9-CM Coding Handbook With Answers, AHA, Faye Brown, RRA, 1991, pages 126-

128.

Illogical Procedure Code Relationships

V0406 VAGINAL HYSTERECTOMY, LAPAROSCOPIC versus OPEN

new 1/1/97

Guideline: When the conversion from a closed procedure to an open procedure occurs, only the code for

the open procedure is assigned. Therefore, the code for the closed procedure with the

endoscopic approach is not assigned.

V0406 Exclusive check (if match, error) - P045

Procedure Table 68.51 Laparoscopic assisted vaginal cholecystectomy

(LAVH) (closed procedure)

Relational Table 68.59 Other vaginal hysterectomy (open procedure)

References: Coding Clinic for ICD-9-CM, AHA, 1st Quarter 1993, page 17; 2nd Quarter 1995, page 11;

Illogical Procedure Code Relationships

V0407 EPISIOTOMY versus OTHER SPECIFIED EPISIOTOMIES

new 1/1/97

Guideline:

Multiple codes should not be assigned when the classification provides a combination code that clearly identifies all the elements documented in the procedural statement. Only the combination code is assigned when the code fully identifies the procedures involved. When the procedural statement includes both the method of delivery and episiotomy, only the combination code should be coded. Read the "Excludes" note under code 73.6.

V0407 Exclusive check (if match, error) - P025		
Procedure Table	73.6	Episiotomy
Relational Table	72.1 72.21 72.31 72.71	Low forceps operation with episiotomy Mid forceps operation with episiotomy High forceps operation with episiotomy Vacuum extraction with episiotomy

References: ICD-9-CM Codebook, Tabular Section, Excludes Notes under Code 73.6.

Coding Clinic for ICD-9-CM, AHA, May-June 1984, page 5.

Coding Clinic for ICD-9-CM, AHA, Jan-Feb 1986, pages 8-10.

ICD-9-CM Coding Handbook with Answers, AHA, Faye Brown, RRA, 1991, pages 12 and 41-45.

ICD-9-CM Coding Handbook with Answers, AHA, Faye Brown, RRA, 1989, pages 12 and 36-41.

Illogical Procedure Code Relationships

V0408 CHOLECYSTECTOMY, LAPAROSCOPIC versus OPEN

new 1/1/97

Guideline: When the conversion from a closed procedure to an open procedure occurs, only the code for

the open procedure is assigned. Therefore, the code for the closed procedure with the

endoscopic approach is not assigned.

.....

V0408 Exclusive check (if match, error) - P024

Procedure Table 51.23 Laparoscopic cholecystectomy (closed procedure)

Relational Table 51.22 Total cholecystectomy (open procedure)

References: Coding Clinic for ICD-9-CM, AHA, 1st Quarter 1993, page 17; 2nd Quarter 1995, page 11.

Illogical Procedure Code Relationships

V0409 APPENDECTOMY, LAPAROSCOPIC versus OPEN

new 1/1/97

Guideline: When the conversion from a closed procedure to an open procedure occurs, only code for the

open procedure is assigned. Therefore, the code for the closed procedure with the endoscopic

approach is not assigned.

V0409 Exclusive check (if match, error) - P026

Procedure Table 47.01 Laparoscopic appendectomy (closed procedure)

Relational Table 47.09 Other appendectomy (open procedure)

References: Coding Clinic for ICD-9-CM, AHA, 1st Quarter 1993, page 17; 2nd Quarter 1995, page 11;

Illogical Procedure Code Relationships

V0410 INCIDENTAL APPENDECTOMY, LAPAROSCOPIC versus OPEN

new 1/1/97

Guideline: When the conversion from a closed procedure to an open procedure occurs, only the code for

the open procedure is assigned. Therefore, the code for the closed procedure with the

endoscopic approach is not assigned.

V0410 Exclusive check (if match, error) - P027

Procedure Table 47.11 Laparoscopic incidental appendectomy

(closed procedure)

Relational Table 47.19 Other incidental appendectomy (open procedure)

References: Coding Clinic for ICD-9-CM, AHA, 1st Quarter 1993, page 17; 2nd Quarter 1995, page 11;

Illogical Procedure Code Relationships

V0411 LYSIS OF PERITONEAL ADHESIONS, LAPAROSCOPIC versus OPEN

new 1/1/97

Guideline: When the conversion from a closed procedure to an open procedure occurs, only the code for

the open procedure is assigned. Therefore, the code for the closed procedure with the

endoscopic approach is not assigned.

V0411 Exclusive check (if match, error) - P028

Procedure Table 54.51 Laparoscopic lysis of peritoneal adhesions

(closed procedure)

Relational Table 54.59 Other lysis of peritoneal adhesions (open procedure)

References: Coding Clinic for ICD-9-CM, AHA, 1st Quarter 1993, page 17; 2nd Quarter 1995, page 11;

Illogical Procedure Code Relationships

V0412 LYSIS OF PERIRENAL OR PERIURETERAL ADHESIONS,

LAPAROSCOPIC versus OPEN

new 1/1/97

<u>Guideline:</u> When the conversion from a closed procedure to an open procedure occurs, only the code for

the open procedure is assigned. Therefore, the code for the closed procedure with the

endoscopic approach is not assigned.

.....

V0412 Exclusive check (if match, error) - P029

Procedure Table 59.03 Laparoscopic lysis of perirenal or periureteral adhesions

(closed procedure)

Relational Table 59.02 Other lysis of perirenal or periureteral adhesions (open

procedure)

References: Coding Clinic for ICD-9-CM, AHA, 1st Quarter 1993, page 17; 2nd Quarter 1995, page 11;

Illogical Procedure Code Relationships

V0413 LYSIS OF PERIVESICAL ADHESIONS, LAPAROSCOPIC versus OPEN

new 1/1/97

Guideline: When the conversion from a closed procedure to an open procedure occurs, only the code for

the open procedure is assigned. Therefore, the code for the closed procedure with the

endoscopic approach is not assigned.

V0413 Exclusive check (if match, error) - P030

Procedure Table 59.12 Laparoscopic lysis of perivesical adhesions

(closed procedure)

Relational Table 59.11 Other lysis of perivesical adhesions (open procedure)

References: Coding Clinic for ICD-9-CM, AHA, 1st Quarter 1993, page 17; 2nd Quarter 1995, page 11;

Illogical Procedure Code Relationships

V0414 OOPHOROTOMY, LAPAROSCOPIC versus OPEN

new 1/1/97

Guideline: When the conversion from a closed procedure to an open procedure occurs, only the code for

the open procedure is assigned. Therefore, the code for the closed procedure with the

endoscopic approach is not assigned.

V0414 Exclusive check (if match, error) - P031

Procedure Table 65.01 Laparoscopic oophorotomy (closed procedure)

Relational Table 65.09 Other oophorotomy (open procedure)

References: Coding Clinic for ICD-9-CM, AHA, 1st Quarter 1993, page 17; 2nd Quarter 1995, page 11;

Illogical Procedure Code Relationships

V0415 OVARIAN BIOPSY, LAPAROSCOPIC versus OPEN

new 1/1/97

Guideline: When the conversion from a closed procedure to an open procedure occurs, only the code for

the open procedure is assigned. Therefore, the code for the closed procedure with the

endoscopic approach is not assigned.

V0415 Exclusive check (if match, error) - P032

Procedure Table 65.13 Laparoscopic biopsy of ovary (closed procedure)

Relational Table 65.12 Other biopsy of ovary (open procedure)

References: Coding Clinic for ICD-9-CM, AHA, 1st Quarter 1993, page 17; 2nd Quarter 1995, page 11;

V0416 OVARIAN CYST MARSUPIALIZATION, LAPAROSCOPIC versus OPEN

new 1/1/97

Guideline: When the conversion from a closed procedure to an open procedure occurs, only the code for

the open procedure is assigned. Therefore, the code for the closed procedure with the

endoscopic approach is not assigned.

V0416 Exclusive check (if match, error) - P033

Procedure Table 65.23 Laparoscopic marsupialization of ovarian cyst

(closed procedure)

Relational Table 65.21 Other marsupialization of ovarian cyst

(open procedure)

.....

References: Coding Clinic for ICD-9-CM, AHA, 1st Quarter 1993, page 17; 2nd Quarter 1995, page 11;

Illogical Procedure Code Relationships

V0417 WEDGE RESECTION OF OVARY, LAPAROSCOPIC versus OPEN

new 1/1/97

Guideline: When the conversion from a closed procedure to an open procedure occurs, only the code for

the open procedure is assigned. Therefore, the code for the closed procedure with the

endoscopic approach is not assigned.

V0417 Exclusive check (if match, error) - P034

Procedure Table 65.24 Laparoscopic wedge resection of ovary

(closed procedure)

Relational Table 65.22 Other wedge resection of ovary (open procedure)

References: Coding Clinic for ICD-9-CM, AHA, 1st Quarter 1993, page 17; 2nd Quarter 1995, page 11;

Illogical Procedure Code Relationships

V0418 UNILATERAL OOPHORECTOMY, LAPAROSCOPIC versus OPEN

new 1/1/97

Guideline: When the conversion from a closed procedure to an open procedure occurs, only the code for

the open procedure is assigned. Therefore, the code for the closed procedure with the

endoscopic approach is not assigned.

V0418 Exclusive check (if match, error) - P035

Procedure Table 65.31 Laparoscopic unilateral oophorectomy

(closed procedure)

Relational Table 65.39 Other unilateral oophorectomy (open procedure)

References: Coding Clinic for ICD-9-CM, AHA, 1st Quarter 1993, page 17; 2nd Quarter 1995, page 11;

Illogical Procedure Code Relationships

V0419 REMOVAL OF BOTH OVARIES, LAPAROSCOPIC versus OPEN

new 1/1/97

Guideline: When the conversion from a closed procedure to an open procedure occurs, only the code for

the open procedure is assigned. Therefore, the code for the closed procedure with the

endoscopic approach is not assigned.

V0419 Exclusive check (if match, error) - P036

Procedure Table 65.53 Laparoscopic removal of both ovaries at same operative

episode (closed procedure)

Relational Table 65.51 Other removal of both ovaries at same operative episode

(open procedure)

References: Coding Clinic for ICD-9-CM, AHA, 1st Quarter 1993, page 17; 2nd Quarter 1995, page 11;

V0420 REMOVAL OF REMAINING OVARY, LAPAROSCOPIC versus OPEN

new 1/1/97

Guideline: When the conversion from a closed procedure to an open procedure occurs, only the code for

the open procedure is assigned. Therefore, the code for the closed procedure with the

endoscopic approach is not assigned.

V0420 Exclusive check (if match, error) - P037

Procedure Table 65.54 Laparoscopic removal of remaining ovary

(closed procedure)

Relational Table 65.52 Other removal of remaining ovary (open procedure)

References: Coding Clinic for ICD-9-CM, AHA, 1st Quarter 1993, page 17; 2nd Quarter 1995, page 11;

V0421 SIMPLE SUTURE OF OVARY, LAPAROSCOPIC versus OPEN

new 1/1/97

Guideline: When the conversion from a closed procedure to an open procedure occurs, only

the code for the open procedure is assigned. Therefore, the code for the closed

procedure with the endoscopic approach is not assigned.

V0421 Exclusive check (if match, error) - P038

Procedure Table 65.74 Laparoscopic simple suture of ovary

(closed procedure)

Relational Table 65.71 Other simple suture of ovary (open procedure)

References: Coding Clinic for ICD-9-CM, AHA, 1st Quarter 1993, page 17; 2nd Quarter 1995,

V0422 OVARY REIMPLANTATION, LAPAROSCOPIC versus OPEN

new 1/1/97

Guideline: When the conversion from a closed procedure to an open procedure occurs, only

the code for the open procedure is assigned. Therefore, the code for the closed

procedure with the endoscopic approach is not assigned.

V0422 Exclusive check (if match, error) - P039

Procedure Table 65.75 Laparoscopic reimplantation of ovary

(closed procedure)

Relational Table 65.72 Other reimplantation of ovary (open procedure)

References: Coding Clinic for ICD-9-CM, AHA, 1st Quarter 1993, page 17; 2nd Quarter 1995,

V0423 UNILATERAL SALPINGO-OOHPHORECTOMY,

LAPAROSCOPIC versus OPEN

new 1/1/97

Guideline: When the conversion from a closed procedure to an open procedure occurs, only

the code for the open procedure is assigned. Therefore, the code for the closed

procedure with the endoscopic approach is not assigned.

V0423 Exclusive check (if match, error) - P040

Procedure Table 65.41 Laparoscopic unilateral salpingo-oophorectomy (closed procedure)

Relational Table 65.49 Other unilateral salpingo-oophorectomy (open procedure)

(open procedure

References: Coding Clinic for ICD-9-CM, AHA, 1st Quarter 1993, page 17; 2nd Quarter 1995,

V0424 REMOVAL OF BOTH OVARIES AND TUBES, LAPAROSCOPIC versus OPEN

new 1/1/97

Guideline: When the conversion from a closed procedure to an open procedure occurs, only

the code for the open procedure is assigned. Therefore, the code for the closed

procedure with the endoscopic approach is not assigned.

V0424 Exclusive check (if match, error) - P041

Procedure Table 65.63 Laparoscopic removal of both ovaries and tubes at same operative episode (closed procedure)

Relational Table 65.61 Other removal of both ovaries and tubes at same operative episode (open procedure)

References: Coding Clinic for ICD-9-CM, AHA, 1st Quarter 1993, page 17; 2nd Quarter 1995,

V0425 REMOVAL OF REMAINING OVARY AND TUBE,

LAPAROSCOPIC versus OPEN

new 1/1/97

Guideline: When the conversion from a closed procedure to an open procedure occurs, only

the code for the open procedure is assigned. Therefore, the code for the closed

procedure with the endoscopic approach is not assigned.

V0425 Exclusive check (if match, error) - P042

Procedure Table 65.64 Laparoscopic removal of remaining ovary and

tube

(closed procedure)

Relational Table 65.62 Other removal of remaining ovary and tube

(open procedure)

.....

References: Coding Clinic for ICD-9-CM, AHA, 1st Quarter 1993, page 17; 2nd Quarter 1995,

V0426 SALPINGO-OOPHOROPLASTY, LAPAROSCOPIC versus OPEN

new 1/1/97

Guideline: When the conversion from a closed procedure to an open procedure occurs, only

the code for the open procedure is assigned. Therefore, the code for the closed

procedure with the endoscopic approach is not assigned.

V0426 Exclusive check (if match, error) - P043

Procedure Table 65.76 Laparoscopic salpingo-oophoroplasty

(closed procedure)

Relational Table 65.73 Other salpingo-oophoroplasty (open procedure)

References: Coding Clinic for ICD-9-CM, AHA, 1st Quarter 1993, page 17; 2nd Quarter 1995,

V0427 LYSIS OF OVARY AND TUBE ADHESIONS,

LAPAROSCOPIC versus OPEN

new 1/1/97

Guideline: When the conversion from a closed procedure to an open procedure occurs, only

the code for the open procedure is assigned. Therefore, the code for the closed

procedure with the endoscopic approach is not assigned.

V0427 Exclusive check (if match, error) - P044

Procedure Table 65.81 Laparoscopic lysis of adhesions of ovary and fallopian tube (closed procedure)

Relational Table 65.89 Other lysis of adhesions of ovary and fallopian tube (open procedure)

tube (open proceas

References: Coding Clinic for ICD-9-CM, AHA, 1st Quarter 1993, page 17; 2nd Quarter 1995,

page 11; 4th Quarter 1996, page 67.

STOP

NEXT V-EDIT IS V0501